

University of Groningen

Ecological forensics

Jouta, Jeltje; Dietz, Maurine W.; Reneerkens, Jeroen; Piersma, Theunis; Rakhimberdiev, Eldar; Hallgrímsson, Gunnar T.; Pen, Ido

Published in:
Methods in ecology and evolution

DOI:
[10.1111/2041-210X.12695](https://doi.org/10.1111/2041-210X.12695)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Jouta, J., Dietz, M. W., Reneerkens, J., Piersma, T., Rakhimberdiev, E., Hallgrímsson, G. T., & Pen, I. (2017). Ecological forensics: Using single point stable isotope values to infer seasonal schedules of animals after two diet switches. *Methods in ecology and evolution*, 8(4), 492-500.
<https://doi.org/10.1111/2041-210X.12695>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Received Date : 06-Oct-2015

Revised Date : 16-Sep-2016

Accepted Date : 10-Oct-2016

Article type : Research Article

Handling editor: Prof. Robert Freckleton

Ecological forensics: Using single point stable isotope values to infer seasonal schedules of animals after two diet switches

Jeltje Jouta^{*a,b}, Maurine W. Dietz^b, Jeroen Reneerkens^b, Theunis Piersma^{a,b}, Eldar Rakhimberdiev^{a,c}, Gunnar T. Hallgrímsson^d, Ido Pen^b

^a*Department of Marine Ecology, NIOZ Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands*

^b*Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands*

^c*Department of Vertebrate Zoology, Lomonosov Moscow State University, 119991, Moscow, Russia*

^d*Institute of Biology, University of Iceland, Sturlugata 7, 101 Reykjavik, Iceland*^{*}*Corresponding author: Jeltje.Jouta@NIOZ.nl*

RUNNING TITLE

Using isotopes to infer seasonal schedules

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/2041-210X.12695

This article is protected by copyright. All rights reserved.

ABSTRACT

1. Animals adjust to seasonal challenges in physical, behavioural and spatial ways. Such adjustments are commonly associated with diet changes that often can be characterized isotopically.
2. We introduce the 'double diet switch model', with which the occurrence and timing of two subsequent diet switches of an individual animal can be traced with a single sample assayed for stable isotopes. We demonstrate the model for Sanderling, *Calidris alba*, a small shorebird that migrates from the Nearctic tundra breeding grounds to the intertidal flats of the Wadden Sea; during this migration some birds may stage in the North Atlantic areas.
3. The 'double diet switch model' successfully predicted the occurrence and timing of two diet switches in 59 Sanderlings captured in the Wadden Sea in July-September. Excluding birds that likely had over-summered at North Atlantic staging areas, the model predicted that Sanderlings departed from the Arctic on 13 July (range: 9-17 July), had a staging duration of 18.6 days in the North Atlantic, and arrived in the Wadden Sea on 1 August (31 July-1 August). The estimated mean Arctic departure dates coincided with the mean hatching date, suggesting that many individuals failed to produce young or left the care to a partner. Estimated mean arrival date matched the main arrival period in the Wadden Sea obtained from observation data. In this study we did not use lipid-free tissues, which may bias model predictions. After correcting for lipid components, the estimated departure date was 11 days later and the staging duration 8.5 days shorter, while arrival date was similar.
4. The 'double diet switch model' successfully identified the occurrence and timing of two subsequent diet switches. The 'double diet switch model will not only apply to switches between three isotopic levels (as in the case study on Sanderling) but also to scenarios where the second switch reverses to the initial isotopic level. Due to this general applicability, the model can be adapted to a wide range of taxa and situations. Foreseeable applications include changes in

habitat and food type, ontogenetic development, or drastic phenotypic changes such as the metamorphosis in insects and amphibians.

KEYWORDS

Calidris alba – dietary change – migration – seasonal patterns – shorebirds – stable isotopes – staging duration – timing – trophic change – Wadden Sea

INTRODUCTION

Animals adjust to seasonal challenges by movements and by physical and behavioural changes (Piersma & van Gils 2011). Quite commonly, these adjustments are associated with diet changes that can be isotopically characterized (Hobson 1999; Caut, Angulo & Courchamp 2009). The accompanying shifts in isotopic value enables researchers to illuminate seasonal phenomena such as migration, metamorphosis, or (temporary) increasing or declining food availability (Phillips & Eldridge 2006; Karasov & Martínez del Rio 2007; Schwemmer et al. 2016). No surprise that 'ecological forensics' is thriving (Dawson & Siegwolf 2011).

Stable isotope analyses can track the occurrence and timing of diet switches based on differences in (1) isotopic values generated by foraging on isotopically distinct food sources and (2) incorporation times of an isotope in distinct consumer tissues (e.g. plasma and red blood cells: Hobson 1999; Klaassen et al. 2010). After a diet switch, the isotopic incorporation of the new diet in a consumer's tissues follows a first order kinetics model, mostly described by an exponential decay function. This model can estimate the time since a single diet switch by using stable isotope values of one, or preferably two, tissue types (Phillips & Eldridge 2006; Klaassen et al. 2010; Oppel & Powell 2010). For animals that change their foraging location or diet more than once over relatively short

time spans, we here describe a ‘double diet switch model’. This model can deal with three successive isotopically distinct diets based on a single assessment of isotopic values in two tissues with distinct turnover rates in one individual and gives estimates of the timing of the two consecutive diet switches.

To demonstrate the functionality of the model, we estimate the timing of post-breeding migration of Sanderlings *Calidris alba* upon their arrival in the Dutch Wadden Sea. After a breeding season in the High Arctic, these long-distance migratory shorebirds depart from the tundra where they fed on terrestrial arthropods (Wirta et al. 2015). Before arrival in the Wadden Sea, where they mainly feed on Brown Shrimp *Crangon crangon* (JR pers. comm.), Sanderlings may or may not make refuelling stops in coastal habitats in the North Atlantic where soft-bodied marine invertebrates comprise the diet (Reneerkens et al. 2009).

METHODS

THE DOUBLE DIET SWITCH MODEL

The isotopic change of body tissues after a diet switch typically follows a first-order kinetic response which is generally well described by a negative exponential function (Tieszen et al. 1983; Phillips & Eldridge 2006; Klaassen et al. 2010). Specifically, consider a focal animal on a diet A, with a corresponding isotope ratio δ_{A1} in tissue 1. If at time $t = 0$ the animal switches from diet A to diet B, then after t_B days on the new diet, its tissue-specific isotope ratio is given by the formula

$$\delta(t_B) = \delta_{B1} + (\delta_{A1} - \delta_{B1})e^{-\lambda_1 t_B}, \quad (1)$$

where δ_{B1} is the characteristic isotope ratio of diet B in tissue 1, and λ_1 is the tissue-specific turnover rate (1/day) of the isotope. Given estimates of δ_{A1} , δ_{B1} and λ_1 , this ‘single diet switch model’ allows

estimation of t_B , the amount of time since the diet switch occurred (Phillips & Eldridge 2006; Klaassen et al. 2010).

Here we expand this ‘single diet switch model’ to one which describes two diet switches: the ‘double diet switch model’. Suppose that at time $t = t_B$, our focal animal switches once again, from diet B to diet C, the latter having characteristic isotopic ratio δ_{C1} in tissue 1. After t_C days on diet C, at time $t = t_B + t_C$, the animal’s isotope ratio is now given by

$$\begin{aligned}\delta(t) &= \delta_{C1} + [\delta(t_B) - \delta_{C1}]e^{-\lambda_1 t_C} \\ &= \delta_{C1} + [\delta_{B1} + (\delta_{A1} - \delta_{B1})e^{-\lambda_1 t_B} - \delta_{C1}]e^{-\lambda_1 t_C},\end{aligned}\tag{2}$$

where we substituted the right-hand side of formula (1) for $\delta(t_B)$ in the first line. Note that this formula is not very useful by itself, since any observed value of $\delta(t)$ within the range spanned by δ_{A1} , δ_{B1} and δ_{C1} is typically consistent with infinitely many combinations of t_B and t_C . However, if a sample is taken simultaneously from a second tissue with a different turnover rate λ_2 , then we have a system of two equations for the two unknowns t_B and t_C :

$$\begin{aligned}\delta_1(t) &= \delta_{C1} + [\delta_{B1} - \delta_{C1} + (\delta_{A1} - \delta_{B1})e^{-\lambda_1 t_B}]e^{-\lambda_1 t_C} \\ \delta_2(t) &= \delta_{C2} + [\delta_{B2} - \delta_{C2} + (\delta_{A2} - \delta_{B2})e^{-\lambda_2 t_B}]e^{-\lambda_2 t_C}\end{aligned}\tag{3}$$

Geometrically, the two equations correspond to two curves in the t_B - t_C plane, and solutions to the two equations occur if and where the curves intersect. As we shall see below, these solutions are precisely the maximum likelihood estimates of t_B and t_C , provided that $\delta_1(t)$ and $\delta_2(t)$ are normally distributed around their predicted values. Solving both equations for t_C gives explicit formulas for the two curves:

$$\begin{aligned}t_C &= \frac{1}{\lambda_1} \ln \left(\frac{\delta_{C1} - \delta_{B1} + (\delta_{B1} - \delta_{A1})e^{-\lambda_1 t_B}}{\delta_{C1} - \delta_1(t)} \right) \\ t_C &= \frac{1}{\lambda_2} \ln \left(\frac{\delta_{C2} - \delta_{B2} + (\delta_{B2} - \delta_{A2})e^{-\lambda_2 t_B}}{\delta_{C2} - \delta_2(t)} \right)\end{aligned}\tag{4}$$

Equating both right-hand-sides yields an equation in t_B , which does not have closed-form solutions, but which may be solved by standard numerical routines. If a solution is found, it can be put back into either of the right-hand sides of (4) to give a corresponding solution for t_C .

Thus, the ‘double diets switch model’ allows estimation of seasonal scheduling of animals with three subsequent diets, such as migrant birds consuming isotopically distinct diets before the start of migration, during a staging episode and after arrival to final destination, respectively, or grizzly bears (*Ursus arctos*) switching temporarily from a diet with mainly whitebark pine (*Pinus albicaulis*) to a diet with mainly elk (*Cervus elaphus*) (Schwartz et al. 2014). The conditions for the use of the ‘double diet switch model’ are presented in table 1. In the next section we describe a statistical method to estimate t_B and t_C .

THE LIKELIHOOD MODEL

We use a maximum likelihood (ML) approach to estimate the parameters t_B and t_C in the nonlinear model (3), given estimates of all other parameters and the measured values of δ_1 and δ_2 . We assume that measurement errors have a normal density:

$$p(\delta_1, \delta_2 | t_B, t_C) = \frac{1}{2\pi\sigma_\delta^2} \exp\left(-\frac{1}{2\sigma_\delta^2} ((\delta_1 - \mu_1)^2 + (\delta_2 - \mu_2)^2)\right). \quad (5)$$

Here σ_δ^2 is the variance, assumed known and identical for both tissues, while μ_1 and μ_2 are the expected values of δ_1 and δ_2 according to model (3):

$$\begin{aligned} \mu_1(t_B, t_C) &= \delta_{C1} + [\delta_{B1} - \delta_{C1} + (\delta_{A1} - \delta_{B1})e^{-\lambda_1 t_B}]e^{-\lambda_1 t_C} \\ \mu_2(t_B, t_C) &= \delta_{C2} + [\delta_{B2} - \delta_{C2} + (\delta_{A2} - \delta_{B2})e^{-\lambda_2 t_B}]e^{-\lambda_2 t_C} \end{aligned} \quad (6)$$

The log-likelihood is then, up to a constant term:

$$\ell(t_B, t_C) = -\frac{1}{2\sigma_\delta^2}((\delta_1 - \mu_1)^2 + (\delta_2 - \mu_2)^2) \quad (7)$$

The score, the partial derivatives of the log-likelihood with respect to both parameters is then given by

$$\begin{aligned} \frac{\partial \ell}{\partial t_B} &= \frac{1}{\sigma_\delta^2} \left((\delta_1 - \mu_1) \frac{\partial \mu_1}{\partial t_B} + (\delta_2 - \mu_2) \frac{\partial \mu_2}{\partial t_B} \right) \\ \frac{\partial \ell}{\partial t_C} &= \frac{1}{\sigma_\delta^2} \left((\delta_1 - \mu_1) \frac{\partial \mu_1}{\partial t_C} + (\delta_2 - \mu_2) \frac{\partial \mu_2}{\partial t_C} \right) \end{aligned} \quad (8)$$

Clearly the score vanishes if $\mu_1 = \delta_1$ and $\mu_2 = \delta_2$, which shows that the ML estimates of t_B and t_C are indeed the solutions to the system of equations (4). We used the function `uniroot` in R version 3.3.0 (R Core Team 2016) to find numerical solutions. All R scripts are available as online appendices to this paper.

The Hessian matrix of second order derivatives, evaluated at the candidate ML estimates, is

$$H = \begin{bmatrix} \frac{\partial^2 \ell}{\partial t_B^2} & \frac{\partial^2 \ell}{\partial t_B \partial t_C} \\ \frac{\partial^2 \ell}{\partial t_B \partial t_C} & \frac{\partial^2 \ell}{\partial t_C^2} \end{bmatrix} = \frac{1}{\sigma_\delta^2} \begin{bmatrix} -\left(\frac{\partial \mu_1}{\partial t_B}\right)^2 - \left(\frac{\partial \mu_2}{\partial t_B}\right)^2 & -\frac{\partial \mu_1}{\partial t_B} \frac{\partial \mu_1}{\partial t_C} - \frac{\partial \mu_2}{\partial t_B} \frac{\partial \mu_2}{\partial t_C} \\ -\frac{\partial \mu_1}{\partial t_B} \frac{\partial \mu_1}{\partial t_C} - \frac{\partial \mu_2}{\partial t_B} \frac{\partial \mu_2}{\partial t_C} & -\left(\frac{\partial \mu_1}{\partial t_C}\right)^2 - \left(\frac{\partial \mu_2}{\partial t_C}\right)^2 \end{bmatrix} \quad (9)$$

The Hessian has two uses here: first, to verify that candidate ML solutions are indeed maxima of the likelihood, and secondly, to provide approximate standard errors for the ML estimates. A local maximum is verified if $\text{tr}(H) = H_{11} + H_{22} < 0$, which is easily seen to be true, and if $\det(H) = H_{11} H_{22} - H_{12} H_{21} > 0$, which is also true since $\det(H) = \left(\frac{\partial \mu_1}{\partial t_B} \frac{\partial \mu_2}{\partial t_C} - \frac{\partial \mu_1}{\partial t_C} \frac{\partial \mu_2}{\partial t_B}\right)^2 > 0$. Approximate standard errors and covariances for the ML estimates \hat{t}_B and \hat{t}_C follow from

$$-H^{-1} \approx \begin{bmatrix} \sigma_{\hat{t}_B}^2 & \sigma_{\hat{t}_B \hat{t}_C} \\ \sigma_{\hat{t}_B \hat{t}_C} & \sigma_{\hat{t}_C}^2 \end{bmatrix} \quad (10)$$

The matrix $-H$ is called the information matrix, since the inverse of information is uncertainty, as quantified by standard errors. To evaluate H we need to evaluate the partial derivatives for tissues $i = 1, 2$:

$$\begin{aligned} \left. \frac{\partial \mu_i}{\partial t_B} \right|_{t_B = \hat{t}_B, t_C = \hat{t}_C} &= -\lambda_i (\delta_{Ai} - \delta_{Bi}) e^{-\lambda_i (\hat{t}_B + \hat{t}_C)} \\ \left. \frac{\partial \mu_i}{\partial t_C} \right|_{t_B = \hat{t}_B, t_C = \hat{t}_C} &= -\lambda_i (\delta_{Bi} - \delta_{Ci} + (\delta_{Ai} - \delta_{Bi}) e^{-\lambda_i \hat{t}_B}) e^{-\lambda_i \hat{t}_C} \end{aligned} \quad (11)$$

Plugging these into (9) clearly shows that the uncertainty about \hat{t}_B and \hat{t}_C increases exponentially with their estimated mean values. Specifically, according to the first equation in (11), information regarding t_B decays exponentially if either t_B or t_C grows large, while according to the second equation information regarding t_C is especially sensitive to large t_C but not t_B values. Thus, unless turnover rates are very low, it is clearly preferable to sample not too long after the second diet switch, nor should the time between diet switches be too long.

We have attempted to take a full Bayesian approach to estimate t_B and t_C , but the maximum likelihood (ML) approach was superior. Simulations indicated (results not shown) that even weakly informative priors produced considerable bias in estimates. The use of flat priors is ruled out for our model since the likelihood does not converge to zero as t_B and t_C go to infinity, rendering the corresponding posterior distribution non integrable.

SENSITIVITY ANALYSIS

The model has 8 parameters: for each tissue $i = 1, 2$ and diet $j = A, B, C$ the equilibrium isotope ratios are denoted by δ_{ij} and turnover rates by λ_i . For the Sanderling data, the diet-and tissue-specific isotope ratios and associated standard deviations were estimated directly from blood and indirectly

from prey items (table 2, supporting information I). No direct information about turnover rates was available for the Sanderling. Instead values for λ_i were predicted on the basis of interspecific allometric regressions, while standard deviations were obtained as averages of intraspecific standard deviations (table S1, supporting information II).

To assess the sensitivity of model predictions to uncertainty in the 8 parameters, for each bird in our dataset we drew 10000 random normal deviates for each of the 6 isotope ratios and for the logarithms of the turnover rates (which must be positive), based on our estimates of mean values and standard deviations. For the isotope ratios we used independent draws, while for turnover rates we allowed for a positive correlation between tissues since it seems plausible that variation in metabolic rate affects turnover rates in the same direction. For each of the draws we attempted to obtain ML estimates for t_B and t_C by solving system (4). When we obtained a candidate solution, we calculated the Hessian to verify it corresponded to a maximum and to estimate standard errors for the parameter estimates. Thus, for each bird we obtained 1000 distributions, one for each successful random draw, which we approximated as a mixture of 10000 gamma distributions to avoid negative values in the tails of the distributions. The mixture was stored as a “posterior distribution” from which we calculated mean values and 89% highest posterior density intervals.

As an alternative to our simulation approach to sensitivity analysis, parameter likelihoods may also be incorporated into an overall likelihood for all model parameters, in addition to t_B and t_C , and corresponding confidence levels calculated. Such an extended likelihood-approach would have to be tailored to the study-specific way the additional parameters were estimated.

THE CASE: TIMING OF SOUTHWARD MIGRATION IN SANDERLING

Using the ‘double diet switch model’, we reconstructed the timing of southward migration by Sanderlings from the tundra breeding grounds (where they ate diet A) and subsequently flew, with or without staging in the North Atlantic (diet B), to the Wadden Sea (diet C). In July-September 2011

and 2012, 65 adult Sanderlings were captured with mist-nets during new moon nights near high-tide roosts in the western Dutch Wadden Sea (53°N, 4-5°E). In addition, 10 adult Sanderlings were caught on their nests in Greenland (Zackenbergl, 74°30'N, 21°00'W) in the second half of June 2009. Blood samples of these latter birds were used to determine the $\delta^{13}\text{C}$ value of red blood cells (RBC) and plasma of birds on the initial diet in the Arctic (diet A; see supporting information I).

Immediately after capture, all 75 Sanderlings were (colour)ringed, weighed and aged based on plumage criteria (Prater, Marchant & Vuorinen 1977), and a blood sample (~300 μL) for stable isotope analysis was drawn from the brachial vein into heparinised capillaries. Note that second calendar year Sanderlings cannot be distinguished from older Sanderling based on their plumage after their first basic moult in spring (Prater, Marchant & Vuorinen 1977; Lemke, Bowler & Reneerkens 2012). Immediately after sampling, the blood was centrifuged in Eppendorf cups in a haematocrit centrifuge (microfuge Sigma 1-13, 6 min on 5000 rpm). Plasma and RBC were pipetted in separate glass vials and stored in a freezer (-20°C) until analysis.

The Sanderling dataset serves all conditions for the 'double diet switch model', as described in Table 1: (a) Stable carbon isotope analysis were performed on plasma and RBC of Sanderlings caught in the Wadden Sea. (b) The $\delta^{13}\text{C}$ values of plasma and RBC of Sanderlings differed between all three locations along the migration route (Table 2). North Atlantic staging areas were assigned based on eight re-sightings of colour-ringed Sanderlings (2007-2014) recorded within the same season of southward migration at both a North Atlantic staging area and the Wadden Sea (Fig. 1). The isotope values of Sanderling's RBC and plasma at locations A and C were obtained from Sanderling blood samples, while the isotope values of RBC and plasma at the North Atlantic staging location (location B) were estimated via prey tissues and a discrimination factor (see supporting information I). (c) The turnover rates for plasma ($\lambda_{\text{plasma}} = 0.303 \pm 0.033$ SD) and RBC ($\lambda_{\text{RBC}} = 0.056 \pm 0.012$ SD) were estimated for an average adult Sanderling (see supporting information II). (d) The tissue sampling dates of all Sanderlings captured in the Wadden Sea were known. (e) There is no indication for non-uniformity in diet between individual Sanderlings under any of the three diets. Besides, it is unlikely

that individual diet specialisation alters the average stable isotope signature of the diet, because we took all important prey species into account, intra-diet variation was within the limits of inter-diet variation, and the consumed prey species differed between the three sites. (f) Samples were collected in the period shortly after the mean arrival period in the Wadden Sea. The ten samples that were collected in late summer, some weeks after the arrival period, indeed showed that the majority of these birds were already adapted to the Wadden Sea diet (Fig. 2).

Figure 2 shows the predictions of the 'double diet switch model' for Sanderlings with different staging durations. The steepness of the slopes of the model predictions increases with turnover rate of the tissue, showing that plasma $\delta^{13}\text{C}$ values (dashed lines) adapt more quickly to the new diet than RBC $\delta^{13}\text{C}$ values (solid lines). The model is based on the combined differences of values for $\delta^{13}\text{C}_{\text{plasma}}$, $\delta^{13}\text{C}_{\text{RBC}}$ and the difference between plasma and RBC isotope values ($\delta^{13}\text{C}_{\text{plasma}}$ minus $\delta^{13}\text{C}_{\text{RBC}}$) over time (t_b and t_c). Therefore, the seasonal schedule of an individual Sanderling can be predicted using a single time point measurement of the stable isotopic value of two tissues. Birds with an Arctic isotopic value in both RBC and plasma are still in equilibrium with the Arctic diet and must have flown directly to the Wadden Sea. Birds with a very short staging period in the North Atlantic staging area and recently arrived in the Wadden Sea will also show a predominantly Arctic signature. Birds with Wadden Sea isotopic values in both RBC and plasma are birds that have been long enough in the Wadden Sea for both tissues to achieve equilibrium with the Wadden Sea diet. We expect that the 'double diet switch model' cannot assign a staging duration to Sanderlings that are already isotopically resident in the Wadden Sea (cf. Hobson 1999). Birds with intermediate values might have been in the Wadden Sea for some time, but not long enough to be in equilibrium with the Wadden Sea diet, and/or may have staged in the North Atlantic region.

Note that migratory flights from the Arctic breeding area in Greenland to the Wadden Sea, which we expect to last approximately two days (65 km/h ground speed for the whole flight of approx. 2850 km; Zwarts et al. 1990), are not taken into account in the model. Although this could

potentially affect the biological interpretation of departure dates from the Arctic, the time in flight is short in comparison with the mean error term of t_b (9.1 days, N=52) . We assumed (1) that a diet switch started upon arrival at a new location and (2) uniform isotopic diets in the three reference areas are representative for the different regions (Arctic, North Atlantic staging areas and Wadden Sea) used by Sanderlings during southward migration to the Wadden Sea. Note also that output dates were reconstructed from termination of 'day of the year' of 2011, since most birds were caught in that year, while the day of the year differs one day between 2011 and 2012.

To evaluate the seasonal schedules of Sanderlings estimated by our 'double diet switch model', we compared our model data with observation data of seasonal schedules of Greenlandic breeding Sanderlings migrating southwards. In 2007-2014, Sanderling nests were annually searched for in northeast Greenland (Reneerkens et al. 2014). Dates of hatch were often exactly known or, in case of clutch predation, estimated based on egg flotation (Hansen et al. 2011). For families found post-hatch, a body mass growth curve based on local data was used to estimate the hatching date. In total we determined hatching dates of 417 clutches and broods (annual range 25-77). The timing of southward migration of Sanderlings was determined based on sightings of individually colour-ringed birds. More than 5600 Sanderlings were individually marked in 12 countries produced over 58,000 unique observations along the East Atlantic flyway collected by us and many volunteers. This dataset was used to extract information of birds sighted in the North Atlantic region and the Wadden Sea within the same season of southward migration.

STABLE ISOTOPE ANALYSIS

All bird plasma, RBC and prey items were stored at -20°C before analysis. The samples were freeze-dried before grinding them with a mortar and pestle. We used a microbalance (Sartorius CP2P) to weigh 0.4 – 0.8 mg of the sample material in 5x8 mm tin capsules. The $\delta^{13}\text{C}$ values were determined with a Thermo Flash 2000 elemental analyser coupled to a Thermo Delta V isotope ratio mass

spectrometer. Isotope values were calibrated to a laboratory acetanilide standard ($\delta^{13}\text{C}$ -26.1 ‰ calibrated on NBS-22) and corrected for blank contribution. 72% of the plasma and RBC samples were analysed in duplicate. The results are reported on the per mill scale with respect to Vienna Pee Dee Belemnite [VPDB]. The replicate error on the standard, acetanilide, ranged between 0.03 and 0.08, using one standard every 4.3 to 7 bird samples.

ELIMINATION OF BIRDS OVERSUMMERING IN THE NORTH ATLANTIC REGION

Our dataset on stable isotope profiles appeared to contain Sanderlings that probably over-summered in the North Atlantic 'staging area' and did not migrate to the Arctic tundra. The estimated staging duration of these individuals was so exceptionally long that if they would have arrived from the Arctic they would have had to depart unrealistically early (as early as 14 May, when Sanderlings are still on northward migration to the Arctic). The 'double diet switch model' cannot eliminate birds that over-summered in the North Atlantic, but simply predicts that these birds have exceptionally long staging durations. In order to eliminate the birds that may have over-summered in the North Atlantic, we excluded birds with a $\delta^{13}\text{C}_{\text{RBC}}$ that fell within or was higher than the $\delta^{13}\text{C}$ of the North Atlantic staging area and also had a $\delta^{13}\text{C}_{\text{plasma}}$ that was still not yet adapted to the Wadden Sea diet (7 birds; see Fig. 3A).

RESULTS

The $\delta^{13}\text{C}$ values of RBC and plasma of Sanderlings caught in the Wadden Sea varied from -24.32 ‰, which is close to a signature of bird's blood in equilibrium with a diet on the Arctic terrestrial arthropods, to -13.5 ‰, which is a signature for bird's blood in equilibrium with the Wadden Sea diet (Fig. 3A). Whereas most birds captured in late summer showed Wadden Sea diet type isotopic values in both RBC and plasma, birds captured in the main arrival period (23 July to 2 August) showed a

variety of patterns ranging from almost purely Arctic signatures, North Atlantic isotopic signatures, intermediate isotopic values, to Wadden Sea diet signatures (Fig. 3A).

Based on the 'double diet switch model' we assessed the individual seasonal schedules of the Sanderlings (Fig. 3). Sanderlings had a wide range of migration strategies with staging periods along North Atlantic coasts ranging from 2.2 to 37.6 days (Fig. 3B). Sanderlings departed from the Arctic on average on 13 July (range: 9-17 July, N=52, Fig. 4), to arrive in the Wadden Sea on 1 August (31 July-1 August, N=52, Fig. 4). When we include the seven birds that over-summered in the North Atlantic staging areas, the mean arrival date remained 1 August (range: 31 July-1 August, N=59, Fig. 4). Departure dates from the Arctic and arrival dates in the Wadden Sea for all individual birds are presented in Fig. 4B.

DISCUSSION

Here we developed a new inferential statistical tool to estimate the timing of movements between distinct habitats on the basis of chemical markers in animal tissues. Ecological forensic problems by their nature are particular and specific, and for this reason we will discuss the Sanderling case before zooming out to the wider range of situations to which our new tool can be applied.

Interestingly, with the help of the 'double diet switch model', we are the first to describe the timing of southward migration of Sanderlings. Our results shows that Sanderlings that spend the summer in the Arctic, as well as those which over-summered in the North Atlantic, arrive simultaneously in the Wadden Sea, matching the main arrival date obtained by observations (Loonstra, Piersma & Reneerkens 2016). As surmised by Reneerkens et al. (2009), the 'double diet switch model' revealed that Sanderlings show large temporal variation in the autumn migration schedules. Contrary to the work of Dietz et al. (2010) who, with the help of a 'single diet switch model' found that Red Knots *Calidris canutus* do not stage in the North Atlantic during southward

Accepted Article

migration, we show that Sanderlings stage for variable lengths of time in the North Atlantic before moving on the Wadden Sea. The mean staging duration in coastal areas between Greenland and the Netherlands of southward migrating Sanderlings was estimated to last 18.6 days. The mean departure date from the Arctic was estimated as 13 July. This coincides with the mean hatching date in northeast Greenland (13 July). The majority of clutches fails due to depredation (Reneerkens et al. 2014) and Sanderlings often leave their partner with the care of eggs (Reneerkens et al. 2011). When clutches are incubated by two adults, one of the partners always leaves the other parent with the chicks, as soon as they hatch (Reneerkens et al. 2014). This would explain the early departures from the Arctic tundra by the majority of assayed birds. The seven individuals that seemed to have over-summered in the North Atlantic were most likely second calendar year birds (Summers, Underhill & Prŷs-Jones 1995). The proportion over-summering Sanderlings in the North Atlantic (12%) is comparable to an earlier study by Lemke, Bowler and Reneerkens (2012) who estimated the percentage of juveniles in a wintering population in Scotland to be 6 – 9%.

At time of our isotope analyses, it was not common practice to use lipid-free tissues. It is clear now that lipids may influence isotopic values substantially, also in blood tissue (e.g. Rode et al. 2016). Specifically, high lipid contents in tissue biases $\delta^{13}\text{C}$ values downwards, while lipid contents may vary between individual and tissue type. Although our case study with Sanderlings clearly demonstrates the applicability of the double diet switch model, the estimated migration schedule may be biased for not using lipid-free tissues. To explore this possible bias, we corrected for lipid contents following the method of Post et al. (2007), who suggested to use C:N ratios of the sampled tissue to correct for lipid contents by adding a correction term to the estimated $\delta^{13}\text{C}$ values, and we reran the model with the 'lipid-free' approximate $\delta^{13}\text{C}$ values (of all tissues, from Sanderlings and prey). Using the 'lipid-free' data, the model did not converge for 14 birds (while all 59 birds converged when using uncorrected values), indicating that corrections were inconsistent with the model. Using the approximated 'lipid-free' data of the remaining birds, resulted in an estimated departure date from the Arctic that was later than when using uncorrected data (24 July [CI 20 - 26

July], rather than 13 July), a shorter estimated staging duration (10.1 days [CI 7.6 - 14.9], rather than 18.6 days), but a similar arrival date in the Wadden Sea (31 July [29 July - 2 August] compared with 1 August) (N=45). The model estimates using the 'lipid-free' data matched better with our expectations on the timing of southward Sanderling migration.

As it is likely that Sanderlings show moderate intraspecific variation, we used distributions of the input parameters rather than the mean values, for two reasons. First, individual dietary preferences cause stable isotopic values to vary slightly among individuals. Moreover, the discrimination factor that may be used to distinguish between diet and consumer may vary between individuals as well (supporting information I; Caut, Angulo & Courchamp 2009). Second, intraspecific variation in turnover rates is rather large and poorly understood (Martínez del Río et al. 2009; Hahn et al. 2012). More accurate information about intraspecific variation in turnover rates is needed for more accurate estimations of individual seasonal scheduling. As the conditions for using the 'double diet switch model' can be met rather easily on the basis of a single time point stable isotope measurement of the target species (Table 1), the 'double diet switch model' allows a relatively simple way to assess seasonal schedules.

We encourage future use of our model for estimation of seasonal schedules of animals and emphasize that other isotopes than carbon can also be used (e.g. nitrogen or sulphur). The 'double diet switch model' might be particularly interesting in deciphering the timing and occurrence of migration in other migratory animals, animals with changes in food availability during a season (e.g. an animal that follows the food peak of different prey species), or in the timing of ontogenetic development of animals (e.g. from egg to juvenile to adult). Although not tested here, the 'double diet switch model' might not be limited to studies with switches between three isotopic levels, i.e. with diet switches from diet A to B to C, but might also be applicable to scenarios where the second switch reverses to the initial isotopic level, so a double diet switches from diet A¹ to B and from B back to A². We call this an 'ABBA switch' (see Fig. 5). An ABBA switch may occur under temporary

changing conditions such as e.g. breeding, drought, frozen foraging surfaces (no access to regular food) or injuries of the animal that restricts regular prey consumption. The ABBA switch could, theoretically, be studied with the regular formula of the ‘double diet switch model’ (see equation 2), where diet A² can be interpreted in the model as diet C. The model is thus generally applicable, and can be adapted to a wide range of taxa and situations in which animals use two or three distinct diets within a short period of time.

ACKNOWLEDGEMENTS

We thank Bernard Spaans, Allert Bijleveld, Anne Dekinga and others for helping us catch birds in the Wadden Sea. We are grateful to the crews of the RV *Navicula* (NIOZ), RV *Stern* (NIOZ) and MS *Phoca* (Dutch ministry of Economic Affairs) for bringing us to the catching location. JR thanks the Zackenberg logistical team at the Department of Bioscience–Roskilde, Aarhus University, for providing logistics at the research station at Zackenberg, northeast Greenland. Obeying the Dutch laws, field work was carried out under animal welfare (DEC) protocol NIOZ-10.04 amendment 1. Funding from World Wildlife Fund (the Netherlands) and INTERACT (project INTERPRED) under the European Community’s Seventh Framework Programme (grant number 262693) to JR and an International Polar Year grant (NWO) to TP and JR, made the fieldwork in Greenland possible. We thank Stefan Schouten and Kevin Donkers for help with the isotope analysis. We thank the reviewers for their valuable contributions that improved our manuscript. This study was carried out within the projects ‘Waddensleutels’ (WF203930, JJ and TP) and ‘Metawad’ (WF209925, JR, ER and TP), both funded by Waddenfonds.

DATA ACCESSIBILITY

Data is deposited in the Dryad repository <http://dx.doi.org/10.5061/dryad.t72b0>.

AUTHOR CONTRIBUTION STATEMENT

JJ, IP, MD and ER conceived the ideas and designed methodology; JJ, JR, GH and MD collected the data; JJ, MD, IP, JR and TP analysed the data; JJ (and MR and IP) led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

REFERENCES

- Caut, S., Angulo, E. & Courchamp, F. (2009) Variation in discrimination factors ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$): the effect of diet isotopic values and applications for diet reconstruction. *Journal of Applied Ecology*, **46**, 443-453, doi: 10.1111/j.1365-2664.2009.01620.x.
- Dawson, T.E. & Siegwolf, R. (2011) *Stable isotopes as indicators of ecological change*. Academic Press.
- Dietz, M.W., Spaans, B., Dekinga, A., Klaassen, M., Korthals, H., van Leeuwen, C. & Piersma, T. (2010) Do Red Knots (*Calidris canutus islandica*) routinely skip iceland during southward migration? *Condor*, **112**, 48-55, doi: 10.1525/cond.2010.090139.
- Hahn, S., Hoyer, B.J., Korthals, H. & Klaassen, M. (2012) From food to offspring down: Tissue-specific discrimination and turn-over of stable isotopes in herbivorous waterbirds and other avian foraging guilds. *Plos One*, **7**, doi: 10.1371/journal.pone.0030242.
- Hobson, K.A. (1999) Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia*, **120**, 314-326.
- Karasov, W.H. & Martínez del Río, C. (2007) *Physiological Ecology: How Animals Process Energy, Nutrients, and Toxins*. Princeton University Press, Princeton.
- Klaassen, M., Piersma, T., Korthals, H., Dekinga, A. & Dietz, M.W. (2010) Single-point isotope measurements in blood cells and plasma to estimate the time since diet switches. *Functional Ecology*, **24**, 796-804, doi: 10.1111/j.1365-2435.2010.01689.x.
- Lemke, H.W., Bowler, J. & Reneerkens, J. (2012) Establishing the right period to estimate juvenile proportions of wintering Sanderlings via telescope scans in western Scotland. *Wader Study Group Bulletin*, **119**, 129-132.
- Loonstra, A.H.J., Piersma, T. & Reneerkens, J. (2016) Staging duration and passage population size of Sanderlings in the western Dutch Wadden Sea. *Ardea*, **104**, 49-61.
- Martínez del Río, C., Wolf, N., Carleton, S.A. & Gannes, L.Z. (2009) Isotopic ecology ten years after a call for more laboratory experiments. *Biological Reviews*, **84**, 91-111, doi: 10.1111/j.1469-185X.2008.00064.x.
- Oppel, S. & Powell, A.N. (2010) Carbon isotope turnover in blood as a measure of arrival time in migratory birds using isotopically distinct environments. *Journal of Ornithology*, **151**, 123-131.
- Phillips, D.L. & Eldridge, P.M. (2006) Estimating the timing of diet shifts using stable isotopes. *Oecologia*, **147**, 195-203, doi: 10.1007/s00442-005-0292-0.
- Piersma, T. & van Gils, J.A. (2011) *The flexible phenotype. A body-centred integration of ecology, physiology, and behaviour*. Oxford University Press, Oxford.
- Post, D.M., Layman, C.A., Arrington, D.A., Takimoto, G., Quattrochi, J. & Montana, C.G. (2007) Getting to the fat of the matter: models, methods and assumptions for dealing with lipids in stable isotope analyses. *Oecologia*, **152**, 179-189, doi: 10.1007/s00442-006-0630-x.

- Prater, T., Marchant, J. & Vuorinen, J. (1977) *Guide to the identification and ageing of Holarctic waders*. British Trust for Ornithology, Tring, UK.
- Reneerkens, J., Benhoussa, A., Boland, H., Collier, M., Grond, K., Günther, K., Hallgrimsson, G.T., Hansen, J., Meissner, W., de Meulenaer, B., Ntiamoa-Baidu, Y., Piersma, T., Poot, M., van Roomen, M., Summers, R.W., Tomkovich, P.S. & Underhill, L.G. (2009) Sanderlings using African-Eurasian flyways: a review of current knowledge. *Wader Study Group Bulletin*, **116**, 2-20.
- Reneerkens, J., Grond, K., Schekkerman, H., Tulp, I. & Piersma, T. (2011) Do uniparental Sanderlings *Calidris alba* increase egg heat input to compensate for low nest attentiveness? *Plos One*, **6**, doi:10.1371/journal.pone.0016834.
- Reneerkens, J., van Veelen, P., van der Velde, M., Luttikhuisen, P. & Piersma, T. (2014) Within-population variation in mating system and parental care patterns in the Sanderling (*Calidris alba*) in northeast Greenland. *Auk: Orthological Advances*, **131**, 235-247, doi: 10.1642/auk-13-247.1.
- Rode, K.D., Stricker, C.A., Erlenbach, J., Robbins, C.T., Cherry, S.G., Newsome, S.D., Cutting, A., Jensen, S., Stenhouse, G., Brooks, M., Hash, A. & Nicassio, N. (2016) Isotopic incorporation and the effects of fasting and dietary lipid content on isotopic discrimination in large carnivorous mammals. *Physiological and Biochemical Zoology*, **89**, 182-197, 10.1086/686490.
- Schwartz, C.C., Fortin, J.K., Teisberg, J.E., Haroldson, M.A., Servheen, C., Robbins, C.T. & Van Manen, F.T. (2014) Body and diet composition of sympatric black and grizzly bears in the Greater Yellowstone ecosystem. *Journal of Wildlife Management*, **78**, 68-78, doi: 10.1002/jwmg.633.
- Schwemmer, P., Voigt, C.C., Corman, A.M., Adler, S. & Garthe, S. (2016) Body mass change and diet switch tracked by stable isotopes indicate time spent at a stopover site during autumn migration in dunlins (*Calidris alpina alpina*). *Journal of Avian Biology*, doi: 10.1111/jav.00873.
- Summers, R.W., Underhill, L.G. & Prÿs-Jones, R.P. (1995) Why do young waders in southern Africa delay their first return migration to the breeding grounds? *Ardea*, **83**, 351-357.
- Tieszen, L.L., Boutton, T.W., Tesdahl, K.G. & Slade, N.A. (1983) Fractionation and turnover of stable carbon isotopes in animal tissues: Implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia*, **57**, 32-37, doi: 10.1007/bf00379558.
- Wirta, H., Vesterinen, E., Hambäck, P., Weingartner, E., Rasmussen, C., Reneerkens, J., Schmidt, N.M., Gilg, O. & Roslin, T. (2015) Exposing the structure of an Arctic food web. *Ecology and Evolution*, **17**, 3842-3856, doi: 10.1002/ece3.1647.
- Zwarts, L., Ens, B.J., Kersten, M. & Piersma, T. (1990) Moults, mass and flight range of waders ready to take off for long-distance migrations. *Ardea*, **78**, 339-364.

LIST OF SUPPORTING INFORMATION

- S1: Estimating diet-specific $\delta^{13}\text{C}$ values
- S2: Estimating $\delta^{13}\text{C}$ turnover rates in Sanderlings
- S3: R-scripts

TABLES & FIGURES

Table 1. An overview of the required conditions for the ‘double diet switch model’ to estimate the timing of two consecutive diet switches

Conditions for using the ‘double diet switch model’:	
a)	Stable isotope analysis (e.g. $\delta^{13}\text{C}$) of individuals of the study species should be measured while on the third diet. This should be done for two tissue types: one with a relatively high turnover rate such as plasma and one with a relatively low turnover rate such as RBC). Both tissues should be sampled at the same moment. Tissue sampling needs to be performed before the individual has reached isotopic adaptation of the new equilibrium of the third diet.
b)	At all three stages (or locations), the stable isotope values of the study species itself or that of its food are known (plus a discrimination factor; but see supporting information I) and sufficiently distinct from each other. Ideally, stable isotope values are known for both separate tissue types at all three stages.
c)	Turnover rates of the two tissues are known for the study species (or can be estimated sufficiently accurately).
d)	Preferably, sampling dates of the tissue types are known. With this information durations of the use of a diet can be transferred to dates instead of number of days.
e)	Diet uniformity among individuals.
f)	The sampling moment is important, since there should not have been enough time to approach equilibration to diet C. Besides, the animal’s staging duration should be shorter than the time to approach the equilibration to diet B.

Table 2. Summary of all general input variables of the ‘double switch model’ to estimates individual schedules in migrating Sanderling. Presented are the $\delta^{13}\text{C}$ values of Sanderling in equilibrium with the diets on the three locations along southward migration (mean \pm SE). The $\delta^{13}\text{C}$ values were calculated in two ways and shown in two columns: obtained from Sanderling blood (True) and a calculated value with help of $\delta^{13}\text{C}$ values of prey and a discrimination factor (Calc.). The results of the two methods did not differ significantly (see t-test in last column and supporting information I). Bold values were used in the model.

Diet	Tissue type	Calc. (prey +DiF) ⁴	N	True (bird blood)	N	t-test
Arctic ¹	plasma			-25.99±0.29 ‰	10	
	RBC			-25.33±0.29 ‰		
Staging area ⁶	plasma	-18.29±0.24 ‰	25	-18.28±0.14 ² ‰	4	t(27)=0.02, p=0.99
	RBC	-17.62±0.24 ‰		-17.94±0.30 ² ‰		t(27)=0.53, p=0.60
Wadden Sea	plasma	-14.56±0.09 ‰	20	-14.54±0.16 ³ ‰	6	t(24)=0.16, p=0.91
	RBC	-13.90±0.09 ‰		-13.94±0.13 ³ ‰		t(24)=0.23, p=0.82
Turnover rate ⁵	Tissue type	Mean	SD			
	plasma	0.303	0.033			
	RBC	0.056	0.012			

¹ Based on blood of Sanderlings caught in northeast Greenland.

² Blood of Sanderlings caught in Wadden Sea in summer with $\delta^{13}\text{C}$ values of plasma and RBC that both represented the staging location ($\delta^{13}\text{C}_{\text{plasma}}$ minus $\delta^{13}\text{C}_{\text{RBC}} < 0.23$ ‰). These birds were suspected to have just arrived in the Wadden Sea after using a staging area somewhere in the North Atlantic.

³ Blood of Sanderlings caught in September in the Wadden Sea with $\delta^{13}\text{C}$ values of plasma and RBC that both represented the Wadden Sea ($\delta^{13}\text{C}_{\text{plasma}}$ minus $\delta^{13}\text{C}_{\text{RBC}} < 0.23$ ‰).

⁴ See supporting information I for details about indirect calculations of the $\delta^{13}\text{C}$ signal of Sanderlings. DiF = discrimination factor.

⁵ See supporting information II for calculation of the turnover rate of $\delta^{13}\text{C}$ in plasma and RBC of Sanderlings.

⁶ North Atlantic staging area









